Fluid Phase Equilibria, 62 (1991) 267–268
Elsevier Science Publishers B.V., Amsterdam

Contents Volume 62 (1991)

Volume 62, Nos. 1–2

Predicting the vapor pressures of pure compounds by using the Wagner equation
A. Vetere (San Donato Milan, Italy) 1– 10

An analytic correlation for the vapour pressure of aqueous and non-aqueous solutions of single and mixed electrolytes. Part II. Application and extension
L.A. Cisternas and E.J. Lam (Antofagasta, Chile) 11– 27

Thermodynamic consistency test of vapor–liquid equilibrium data—alcohol–hydrocarbon systems
H.M. Moon, K. Ochi and K. Kojima (Tokyo, Japan) 29– 40

Calculation of vapour–liquid equilibrium data for binary chlorofluorocarbon mixtures using the Peng–Robinson equation of state
S.I. Abu-Eishah (Irbid, Jordan) 41– 52

Phase diagrams of new aqueous phase systems composed of aliphatic alcohols, salts and water
A. Greve and M.-R. Kula (Jülich, Germany) 53– 63

Thermodynamic properties of two alternative refrigerants: 1,1,-dichloro-2,2,2-trifluoroethane (R123) and 1,1,1,2-tetrafluoroethane (R134a)
G. Morrison and D.K. Ward (Gaithersburg, MD, U.S.A.) 65– 86

Critical loci for binary chloroalkane–n-alkane mixtures. III. 1,1-Dichloroethane with C_3–C_9 n-alkanes
F. García-Sánchez and A. Trejo (Mexico, Mexico) 87– 95

Liquid–liquid equilibria for the systems: water/1-pentanol/acetic acid and water/1-hexanol/acetic acid
M.M. Esquível and M.G. Bernardo-Gil (Lisbon, Portugal) 97–107

Excess enthalpies of mixing of acetonitrile with chlorinated ethanes and ethenes
K.N. Surendranath and A. Krishnaiah (Tirupati, India) and M. Ramakrishna (Hyderabad, India) 109–114

PVT data from Burnett and refractive index measurements for the nitrogen–hydrogen system from 270 to 353 K and pressures to 30 MPa
M. Jaeschke and H.M. Hinze (Dorsten, Germany) and H.J. Achtermann and G. Magnus (Hannover, Germany) 115–139

Solid–liquid phase equilibria in $Cu(O_2CCH_3)_2$ + pyridine derivative + chlorobenzene systems
I. Uruska and M. Szpakowska (Gdańsk, Poland) 141–152

Gas–liquid critical temperatures of binary mixtures containing a low molecular weight polydimethylsiloxane or hexaethyldisiloxane
G. Christou, C. Tran and C. Young (Parkville, Vic., Australia) 153–162

Isobaric vapor–liquid equilibrium data for furfural with chlorinated hydrocarbons
J.L. Cabezas and S. Beltrán (Burgos, Spain) and J. Coca (Oviedo, Spain) 163–172

Volume 62, No. 3

A simple relation to predict or to correlate the excess functions of multicomponent mixtures
C.-A. Hwang, J.C. Holste and K.R. Hall (College Station, TX, U.S.A.) and G.A. Mansoori (Chicago, IL, U.S.A.) 173–189

New strategy for phase equilibrium and critical point calculations by thermodynamic energy analysis. Part I. Stability analysis and flash
N.R. Nagarajan and A.S. Cullick (Dallas, TX, U.S.A.) and A. Griewank (Argonne, IL, U.S.A.) 191–210

New strategy for phase equilibrium and critical point calculations by thermodynamic energy analysis. Part II. Critical point calculations
N.R. Nagarajan and A.S. Cullick (Dallas, TX, U.S.A.) and A. Griewank (Argonne, IL, U.S.A.) 211–223

Supercritical phase behaviour: description of the solubility of binary solid mixtures in a supercritical solvent with a lattice gas model
P. Balbuena, E. Campanella and L. Gribaudo (Santa Fe, Argentina) 225–238

P–x measurements for ethanol–n-heptane–isobutanol at 303.15 K
V.R. Bhethanabotla and S.W. Campbell (Tampa, FL, U.S.A.) 239–258

Short Communication

Excess volumes of binary systems formed by a pyridine base and an n-alkane in terms of an association model. Part II. β and γ picolines with C_6–C_{10} n-alkanes
A.J. Treszczanowicz and G.C. Benson (Ottawa, Ont., Canada) and T. Kasprzycka-Guttman (Warsaw, Poland) 259–264

Author Index 265

Contents of Volume 62 (1991) 267–268